

## Blut und Blutprodukte (Teil 1 und 2):

# Indikation zur prähospitalen Gabe

## Literatur:

- Pfeifer R et al. (2009) Patterns of mortality and causes of death in polytrauma patients – has anything changed? *Injury* 2009 40: 907-911.
- Cothren CC, Moore EE, Hedegaard HB, Meng K (2007) Epidemiology of Urban Trauma Deaths: A Comprehensive Reassessment 10 Years Later. *World J Surg* 31: 1507-1511.
- Buschmann C et al. (2015) Vermeidbare Todesfälle nach Trauma. *Notfall Rettungsmed* 18 (4): 316-324.
- Davenport RA, Brohi K (2015) Cause of trauma-induced coagulopathy. *Curr Opin Anaesthesiol* 28: 1-8.
- Spahn DR, Bouillon B, Cerny V et al. (2019) The European guideline on management of major bleeding and coagulopathy following trauma. 5th Ed. *Critical Care* 23: 98.
- National Blood Service (2013) Guidelines for the Blood Transfusion Services in the UK. 8th Ed. The Stationery Office, London.
- Rijnhout Tim WH et al. (2019) Is prehospital blood transfusion effective and safe in haemorrhagic trauma patients? A systematic review and meta-analysis. *Injury* 50 (5): 1017-1027.
- Guerado E, Medina A, Mata MI et al. (2016) Protocols for massive blood transfusion: when and why, and potential complications. *Eur J Trauma Emerg Surg* 42: 283-295.
- Schlump CJ, Schöchl H (2017) The role of fibrinogen in trauma-induced coagulopathy. *Hämostaseologie* 34: 29-39.
- Egidi MG, D'Alessandro A, Mandarello G, Zolla L (2010) Troubleshooting in platelet storage temperature and new perspectives through proteomics. *Blood Transfusion* 8: s73-81.
- Kozar RA et al. (2011) Plasma restoration of endothelial glycoadherent model of hemorrhagic shock. *Anesth Analg* 112: 1289-1295.
- Spahn DR, Bouillon B, Cerny V et al. (2013) Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Crit Care* 17: R76.
- Ducloy-Bouthors AS, Jude B, Duhamel A et al. (2011) High-dose tranexamic acid reduces blood loss in post-partum haemorrhage. *Crit Care* 15: R117.
- Lin Z et al. (2016) Tranexamic acid-associated seizures: A meta-analysis. *Seizure* 36: 70-73.
- Goobie SM (2017) Tranexamic acid: still far to go. *Br J Anaesth* 118 (3): 293-295.
- Fachinformation Pfizer Cyklokapron, Stand: Juni 2016
- Rehn M et al. (2019) Effect of prehospital red blood cell transfusion on mortality and time of death in civilian trauma patient. *Shock* 51 (3): 284-288.
- Stellungnahme der IAKH zum prähospitalen Einsatz von Blutprodukten, Januar 2020.
- Wheeler A et al. (2020) Development of prehospital assessment findings associated with massive transfusion. *Transfusion* 60: 6.
- Oyeni BT et al. (2017) Trends in 1029 trauma deaths at a level 1 trauma center: impact of a bleeding control bundle of care. *Injury* 48 (1): 5-12.
- Marik PE et al. (2008) Efficacy of red blood cell transfusion in the critically ill: a systematic review of the literature. *Crit Care Med* 36: 2667-2674.
- van Turenhout et al. (2020) Pre-hospital transfusion of red blood cells. Part 1: A scoping review of current practice and transfusion triggers. *Transfus Med*. DOI: 10.1111/tme.12667
- Thies KC, Truhlář A, Keene D et al. (2020) Pre-hospital blood transfusion – an ESA survey of European practice. *Scand J Trauma Resusc Emerg Med* 28: 79.
- Hilbert-Carius P et al. (2016) S3-Leitlinie Polytrauma/Schwerverletzten Behandlung – Update.
- Shackelford SA, del Junco DJ, Powell-Dunford N et al. (2017) Association of Prehospital Blood Product Transfusion During Medical Evacuation of Combat Casualties in Afghanistan With Acute and 30-Day Survival. *JAMA* 318: 1581-1511.
- Sperry JL et al. (2018) Prehospital Plasma during Air Medical Transport in Trauma Patients at Risk for Hemorrhagic Shock. *N Engl J Med* 379: 315-326.
- Inaba K, Karamanos E, Lustenberger T et al. (2013) Impact of fibrinogen levels on outcomes after acute injury in patients requiring a massive transfusion. *J Am Coll Surg* (2): 290-297.
- Rourke C, Curry N, Khan S, et al. (2012) Fibrinogen levels during trauma hemorrhage, response to replacement therapy, and association with patient outcomes. *Jf Thrombosis and Haemostasis* 10: 1342-1351.
- Garrigue D, Godier A, Glacet A et al. (2017) French lyophilized plasma versus fresh frozen plasma for the initial management of trauma-induced coagulopathy: a randomized open-label trial. *Journal of Thrombosis and Haemostasis* 16 (3): 481-489.
- COMBAT Study; Moore HB et al. (2018) Plasma-first resuscitation to treat haemorrhagic shock during emergency ground transportation in an urban area: a randomised trial. *Lancet* 392 (10144): 283-291.
- Sperry JL et al. (2018) Prehospital Plasma during Air Medical Transport in Trauma Patients at Risk for Hemorrhagic Shock. *N Engl J Med* 379: 315-326.
- Pusateri AE et al. (2020) Association of Prehospital Plasma Transfusion with Survival in Trauma Patients with Hemorrhagic Shock when Transport Times are Longer than 20 Minutes. A Post Hoc Analysis of the PAMPer and COMBAT Clinical Trials. *JAMA Surg* 155 (2): e195085.
- Stanworth SJ, Brunskill SJ, Hyde CJ, McClelland DB, Murphy MF (2004) Is fresh frozen plasma clinically effective? A systematic review of randomized controlled trials. *Br J Haematol* 126: 139-52.
- Labarinas S et al. (2013) Plasma in the PICU: why and when should we transfuse? *Annals of Intensive Care* 3: 16.
- Roback J et al. (2010) Evidence-based practice guidelines for plasma transfusion. *Transfusion* 50: 1227-1239.
- Neeki MM et al. (2018) Tranexamic Acid in Civilian Trauma Care in the California Prehospital Antifibrinolytic Therapy Study. *West J Emerg Med* 19 (6): 977-986.
- Guyette FX et al. (2020) Tranexamic Acid During Prehospital Transport in Patients at Risk for Hemorrhage After Injury. *JAMA Surg* 156 (1): 11-20.
- Drew B et al. (2020) The Use of Tranexamic Acid in Tactical Combat Casualty Care. *J Spec Oper Med* 3: 36-43.
- Simranjeet B et al. (2019) Mortality and Thrombosis in Injured Adults Receiving Tranexamic Acid in the Post-CRASH-2 Era. *West J Emerg Med* 20 (3): 443-453.
- Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e.V. (AWMF) (2016) Überarbeitete S3-Leitlinie Polytrauma/Schwerverletzten-Behandlung 012-019. [www.awmf.org/leitlinien/detail/II](http://www.awmf.org/leitlinien/detail/II)
- Kozek-Langenecker SA et al. (2017) Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 34: 332-395.
- Rossaint R, Bouillon B et al. (2016) The European guideline on management of major bleeding and coagulopathy following trauma. 4th Ed. *CritCare* 20: 100.
- Schlump CJ et al. (2016) Fibrinogen levels in trauma patients during the first seven days after fibrinogen concentrate therapy: a retrospective study. *Scand J Trauma Resusc Emerg Med* 24: 29.

Autor:

**Dr. Hanifi Dolu, FTLA**  
 Facharzt  
 Klinik für Anästhesiologie, Intensivmedizin, Notfallmedizin u. Schmerztherapie  
 Bundeswehrkrankenhaus Hamburg  
 22049 Hamburg  
[hanifidolu@bundeswehr.org](mailto:hanifidolu@bundeswehr.org)

43. Inaba K et al. (2013) Impact of fibrinogen levels on outcomes after acute injury in patients requiring a massive transfusion. *J Am Coll Surg* 216 (2): 290-297.
44. Heindl B (2012) Evidenzbasierte Therapie mit FFP und Faktorenkonzentraten. Präsentation DIVI.
45. Bundesärztekammer (2020) Querschnittsleitlinie zur Therapie mit Blutkomponenten und Plasmaderivaten. 4. Aufl. [www.bundesaerztekammer.de/fileadmin/user\\_upload/downloads/pdf-Ordner/MuE/Querschnitts-Leitlinien\\_BAEK\\_zur\\_Therapie\\_mit\\_Blutkomponenten\\_und\\_Plasmaderivaten-Gesamtnovelle\\_2020.pdf](http://www.bundesaerztekammer.de/fileadmin/user_upload/downloads/pdf-Ordner/MuE/Querschnitts-Leitlinien_BAEK_zur_Therapie_mit_Blutkomponenten_und_Plasmaderivaten-Gesamtnovelle_2020.pdf) (Abruf: 3. Mai 2021).
46. Goobie SM, Shander A (2020) One Size does not Fit all in Treating Massive Hemorrhage. *Anesthesia & Analgesia* 131: 480-482.
47. Lendrum RA, Kotze JP, Lockey DJ, Weaver AE (2013) Case studies in prehospital care from London HEMS: pre-hospital administration of prothrombin complex concentrate to the head-injured patient. *Emerg Med J* 30: 247-248.
48. Zeeshan M et al. (2019) Four-factor prothrombin complex concentrate is associated with improved survival in trauma-related hemorrhage: A nationwide propensity-matched analysis. *J Trauma Acute Care Surg* 87 (2): 274-281.
49. Schöchl H et al. (2014) Endogenous thrombin potential following hemostatic therapy with 4-factor prothrombin complex concentrate: a 7-day observational study of trauma patients. *Crit Care* 18: R147.
50. Hennig A (2016) Nachweismethoden und Normwerte für Isoagglutinine beim Menschen. Dissertation an der Ludwig-Maximilian Universität München.
51. Nicholas Warner et al. (2018) Military Prehospital Use of Low Titer Group O Whole Blood. *J Spec Oper Med* 18 (1): 15-18.
52. Martinaud C et al. (2018) STORHM Study: French Whole Blood Trial in Traumatic Massive hemorrhage. Department of Clinical Operations French Military Blood Institute, <https://rdcr.org/wp-content/uploads/2018/07/Martinaud-T-STORHM-Martinaud-JUN-2018.pdf> (Abruf: 3. Mai 2021).
53. Strandenes G et al. (2014) Low Titer Group O Whole Blood in Emergency Situations. *Shock* 41: 70-75.
54. Joint Theater Trauma System Clinical Practice Guideline 2012. [www.usaisr.amedd.army.mil/assets/cpgs/Fresh\\_Whole\\_Blood\\_Transfusion\\_24\\_Oct\\_12.pdf](http://www.usaisr.amedd.army.mil/assets/cpgs/Fresh_Whole_Blood_Transfusion_24_Oct_12.pdf) (Abruf: 3. Mai 2021).
55. Food and Drug Administration (2013) Circular of Information: for the use of Human Blood and Blood Components: AABB. [www.aabb.org/resources/bct/Documents/coi0413.pdf](http://www.aabb.org/resources/bct/Documents/coi0413.pdf) (Abruf: 14. April 2021).
56. Tanabe K (2007) Interinstitutional variation in the measurement of anti-A/B antibodies: the Japanese ABO-Incompatible Transplantation Committee survey. *Transplantation* 84 (12) (Suppl.): S13-16.
57. Kumlien G, Wilpert J, Safwenberg J, Tyden G (2007) Comparing the tube and gel techniques for ABO antibody titration, as performed in three European centers. *Transplantation* 84 (12 Suppl): S17-19.
58. AuBuchon JP et al. (2008) Reducing the variation in performance of antibody titrations. *J Lab Med* 132 (7): 1194-1201.
59. Cooling LL, Downs TA, Butch SH, Davenport RD (2008) Anti-A and anti-B titers in pooled group O platelets are comparable to apheresis platelets. *Transfusion* 48 (10): 2106-2113.
60. Win N (2011) NHS blood and transplant: high titre anti-A/B testing of donors within NHS Blood and Transplant (NHSBT). <http://hospital.blood.co.uk/library/pdf/INF178.pdf> (Abruf: 14. April 2021).
61. Baker J (2019) Joint Trauma System Clinical Practice Guideline (JTS CPG) Austere Resuscitative and Surgical Care (ARSC) (CPG ID:76). *Journal of special operations medicine* 20: 25-42.
62. Ponschab M et al. (2015) Haemostatic profile of reconstituted blood in a proposed 1:1:1 ratio of packed red blood cells, platelet concentrate and four different plasma preparations. *Anaesthesia* 70: 528-536.
63. Armand R, Hess JR (2003) Treating coagulopathy in trauma patients. *Transfus Med Rev* 17 (3): 223-231.
64. Spinella P (2008) Warm fresh whole blood transfusion for severe hemorrhage: U.S. Military and potential civilian applications. *Crit Care Med* 36 (7 Suppl): S340-345.
65. Berseus O et al. (2013) Risks of hemolysis due to anti-A and anti-B caused by the transfusion of blood or blood components containing ABO-incompatible plasma. *Transfusion* 53 (Suppl 1): 114S-123S.
66. Behrendt KP (2003) Orator: Leitfaden der Feldchirurgie im Bewegungskrieg, S. 40 und Bürkle de la Camp: Erfahrungsbericht, Meran, Juni 1945, in BA-MA, RH 12-23, H 20/461, S. 7 sowie Sammelbericht Nr. 3, Berlin, Januar 1943, in BA-MA, RH 12-23, H 20/810, S. 1 f.; Dissertation an der Medizinischen Fakultät der Albert-Ludwigs-Universität Freiburg im Breisgau.
67. Spinella PC et al (2007) The risks associated with fresh whole blood and RBS transfusions in a combat support hospital. *Crit Care Med* 35: 2576-2581.
68. Hakre S et al. (2011) Transfusion-transmissible viral infections among US military recipients of whole blood and platelets during Operation Enduring Freedom and Operation Iraqi Freedom. *Transfusion* 51: 473-485.
69. Reed W, Tzong-Hae L, Norris PJ et al. (2007) Transfusion-associated microchimerism: a new complication of blood transfusions in severely injured patients. *Semin Hematol* 44 (1): 24-31.
70. Hoitz J, Kerschowski J (2020) Warm blood transfusions in the medical service for military personnel: an old technique still relevant today?! In: Willy C (Hrsg.) *Worldwide Deployment – Bundeswehr Medical Service 2020*. Beta Verlag, Bonn.
71. Beekley AC (2007) Lessons learned from modern military surgery. *Surg Clin N Am* 87: 157-184.
72. Tom Woolley et al. (2016) NATO Blood Panel perspectives on changes to military prehospital resuscitation policies: current and future practice: *Transfusion* 56 (Suppl 2): S217-223.
73. Sicard B et al. (2016) Bleeding management in remote environment: the use of fresh whole blood transfusion and lyophilised plasma. *Int Marit Health* 67 (2): 79-82.
74. Sicard B et al. (2016) Acute gastrointestinal haemorrhage on board a cruise ship in the Antarctic Peninsula. *Int Marit Health* 67 (4): 223-226.
75. Jenkins D et al. (2014) Implementation and execution of civilian remote damage control resuscitation Programs. *Shock* 41 (Suppl 1): 84-89.
76. Vollmuth R et al. (2015) Wiedervereinigung und Sanitätsdienst. Betrachtungen zu Kontinuität und Wandel. Vorträge des 2. Wehrmedizinischen Symposiums vom 10. November 2010. Beta Verlag, Bonn.
77. Lier H (2017) Stop the bleeding: Gerinnungstherapie bei (Massiv-)Blutungen in der Notaufnahme. Vortrag am Bundeswehrkrankenhaus Hamburg am 27. März 2017. [http://news-papers.eu/wp-content/uploads/2016/12/Lier\\_BlutungZNA\\_Leipzig2016\\_1.pdf](http://news-papers.eu/wp-content/uploads/2016/12/Lier_BlutungZNA_Leipzig2016_1.pdf) (Abruf: 3. Mai 2021).
78. Mabry RL, Holcomb JB, Baker AM et al. (2000) United States Army Rangers in Somalia: an analysis of combat casualties on an urban battlefield. *J Trauma* 49 (3): 515-528.